### **Revised Oswestry Low Back Pain Disability Questionnaire**

From N. Hudson, K. Tome-Nicholson, A. Breen; 1989 rev. 09/11/92

#### Name

Date

# Please mark the ONE choice from EACH group that best describes you. PAIN INTENSITY

- $\Box$ A. The pain comes and goes and is very mild.
- **D**B. The pain is mild and does not vary much.
- $\Box$ C. The pain comes and goes and is moderate.
- D. The pain is moderate and does not vary much.
- $\Box$ E. The pain comes and goes and is severe.
- $\Box$ F. The pain is severe and does not vary much.

#### PERSONAL CARE

- □A. I would not have to change my way of washing or dressing in order to avoid pain.
- **D**B. I do not normally change my way of washing or dressing even though it causes some pain.
- **C**. Washing and dressing increases the pain, but I manage not to change my way of doing it.
- D. Washing and dressing increases the pain and I find it necessary to change my way of doing it.
- **D**E. Because of the pain, I am unable to do some washing and dressing without help.
- **□**F. Because of the pain, I am unable to do any washing or dressing without help.

#### LIFTING

- **A**. I can lift heavy weights without extra pain.
- **D**B. I can lift heavy weights, but it causes extra pain.
- **D**C. Pain prevents me from lifting heavy weights off the floor.
- D. Pain prevents me from lifting heavy weights off the floor, but I can manage if they are conveniently positioned, e.g., on a table.
- □E. Pain prevents me from lifting heavy weights, but I can manage light to medium weights if they are conveniently positioned.
- $\Box$ F. I can only lift very light weights, at the most.

#### WALKING

- **D**A. Pain does not prevent me from walking any distance.
- **D**B. Pain prevents me from walking more than one mile.
- $\Box$ C. Pain prevents me from walking more than  $\frac{1}{2}$  mile.
- D. Pain prevents me from walking more than 1/4 mile.
- $\Box$ E. I can only walk while using a cane or on crutches.
- $\Box$ F. I am in bed most of the time and have to crawl to the toilet.

#### SITTING

- $\Box$ A. I can sit in any chair as long as I like without pain.
- **B**. I can only sit in my favorite chair as long as I like.
- **D**C. Pain prevents me from sitting more than one hour.
- **D**. Pain prevents me from sitting more than  $\frac{1}{2}$  hour.
- $\Box$ E. Pain prevents me from sitting more than ten minutes.
- $\Box$ F. Pain prevents me from sitting at all.



#### STANDING

- **D**A. I can stand as long as I want without pain.
- **D**B. I have some pain while standing, but it does not increase with time.
- **C**. I cannot stand for longer than one hour without increasing pain.
- $\Box$ D. I cannot stand for longer than  $\frac{1}{2}$  hour without increasing pain.
- **D**E. I cannot stand for longer than ten minutes without increasing pain.
- **□**F. I avoid standing, because it increases the pain straight away.

#### SLEEPING

- $\Box$ A. I get no pain in bed.
- **D**B. I get pain in bed, but it does not prevent me from sleeping well.
- **C**. Because of pain, my normal night's sleep is reduced by less than one-quarter.
- D. Because of pain, my normal night's sleep is reduced by less than one-half.
- **D**E. Because of pain, my normal night's sleep is reduced by less than three-quarters.
- $\Box$ F. Pain prevents me from sleeping at all.

#### SOCIAL LIFE

- $\Box$ A. My social life is normal and gives me no pain.
- $\Box$ B. My social life is normal, but increases the degree of my pain.
- □C. Pain has no significant effect on my social life apart from limiting my more energetic interests, e.g., dancing, etc.
- D. Pain has restricted my social life and I do not go out very often.
- $\Box$ E. Pain has restricted my social life to my home.
- $\Box$ F. I have hardly any social life because of the pain.

#### TRAVELING

- $\Box$ A. I get no pain while traveling.
- **D**B. I get some pain while traveling, but none of my usual forms of travel make it any worse.
- □C. I get extra pain while traveling, but it does not compel me to seek alternative forms of travel.
- D. I get extra pain while traveling which compels me to seek alternative forms of travel.
- $\Box$ E. Pain restricts all forms of travel.
- **□**F. Pain prevents all forms of travel except that done lying down.

#### **CHANGING DEGREE OF PAIN**

- **D**A. My pain is rapidly getting better.
- **D**B. My pain fluctuates, but overall is definitely getting better.
- **D**C. My pain seems to be getting better, but improvement is slow at present.
- D. My pain is neither getting better nor worse.
- **D**E. My pain is gradually worsening.
- **□**F. My pain is rapidly worsening.

#### **Patient Signature**

Date



# What's in a number?

How to rate your symptoms, 1 - 10

## Please circle the best that describes your pain

**10** - Your pain is intense, constant, greatly restricts your activities, and it is impossible to go more than 5 minutes without awareness of the pain.

9 - Same as above, but you can forget about the pain up to 15 minutes at a time.

**8** - The pain is significant, moderately intense at times, but not constant. Most activities are affected, you think about it once or twice an hour.

7 - Same as above, except that the pain is never intense.

**6** - The pain is moderate yet too frequent to ignore. Some activities are affected. Hours can go by without being aware of the pain.

5 - Same as above, except that almost no activities are affected.

**4** - The pain is little more than a nuisance, and you go through the whole day frequently aware, but never affected by it.

**3** - Same as above, except that the awareness of the pain may be absent for a whole day at a time.

2 - At its worst, the pain is best described as "a little uncomfortable". Days can go by without being aware of it.

1 - Same as above, except that the symptom does not recur more frequently than once a week.

©1995 The Growth Institute Inc.

Name \_

\_\_\_\_\_Date \_\_\_\_\_

